

Molecular Detection of Human Papilloma Virus (Type 16,18) Using PCR and Its Frequency in Patients with Cervical Cancer in Iranian Women

Zahra Shahi ^{1,2}, Mohammad Amin Edalatmanesh ^{2*}, Babak kheirkhah ³

1. Department of Biology, College of Sciences, Fars Science and Research Branch, Islamic Azad University, Fars, Iran
2. Department of Biology, College of Sciences, Shiraz Branch, Islamic Azad University, Shiraz, Iran
3. Department of Microbiology, College of Sciences, Kerman Branch, Islamic Azad University, Kerman, Iran



Article Info

doi [10.30699/jogcr.5.3.110](https://doi.org/10.30699/jogcr.5.3.110)

Received: 2020/09/20;

Accepted: 2020/10/25;

Published Online: 02 Dec 2020;

Use your device to scan and read the article online



Corresponding Information:

Mohammad Amin Edalatmanesh,
Department of Biology, College of Sciences,
Shiraz Branch, Islamic Azad University,
Shiraz, Iran
Email:
amin.edalatmanesh@gmail.com

ABSTRACT

Background & Objective: Cervical cancer is the fourth most common cancer in women worldwide. HPV is a heterogeneous virus, and a persistent high-risk HPV infection contributes to cancer development. This study aims to determine the relative frequency of HPV genotypes in Kerman, Iran.

Materials & Methods: This cross-sectional study was conducted on 50 women with Pap smear samples, referring to a general laboratory in Kerman, Iran. Detecting two HPV genotypes was carried out using Multiplex Polymerase Chain Reaction (PCR) technique. The sociodemographic survey was conducted for each HPV-positive woman.

Results: Out of 50 cervical cancer patients, 33 women (66%) were HPV 16, 18-positive. HPV 16 (38%) was the most rampant type, followed by HPV 18 (14%) and HPV 16, 18 (14%).

Conclusion: Studying the prevalence of HPV genotypes and their respective risk factors can lead to efficient vaccine development against the virus in each geographical area. It can also be beneficial for illness management and high sensitivity diagnosis of cervical intraepithelial neoplasia.

Keywords: Cervical Cancer, HPV, Kerman, PCR



Copyright © 2020. This is an original open-access article distributed under the terms of the Creative Commons Attribution-noncommercial 4.0 International License which permits copy and redistribution of the material just in noncommercial usages with proper citation.

Introduction

Cancer is the second most common cause of death in developed countries and the third most common cause in developing countries, including Iran (1).

According to the latest epidemiological studies in Iran, cancer is the third leading reason of death following cardiovascular illness and unintentional events. More than 30,000 people die of this disease in Iran annually and it is estimated that more than 70,000 new cases of cancer occur annually in the country (1).

Cancer prevention, early diagnosis and timely treatment are effective in reducing the mortality rate. Infectious agents cause 15-20% of cancers worldwide and 70% of cancers in Eastern Mediterranean countries (2).

With more than 100 genotypes and worldwide contagion, human papillomaviruses (HPVs) are in the *papillomaviridae* family with a circular protein structure and an uncoated ds DNA genome. Papillomaviruses cause a variety of malignant and benign tumors in the human skin and mucous membranes (3). HPV-related diseases and cancers in humans include cancers of the

cervix, skin, larynx, head and neck, breast, genital warts, and conjunctivitis of the eyes and mucous membranes (4, 5).

HPVs are found in 99% of cervical cancer cases with HPV16 and HPV18 being the most prevalent genotypes observed in 70% of cervical carcinomas (6).

Cervical cancer is a malignant illness among women globally. More than 80% of cervical cancer deaths occur in developing countries (7, 8). This study aims to detect the frequency of HPV16, 18 in women with cervical cancer in the Kerman, Iran.

Materials and Methods

Sample separation

In this experimental study, 50 tissue samples related to malignant cervical lesions of different patients were collected from the pathology laboratory of Kerman province (Pasargad laboratory, Iran).

DNA extraction

Virus DNA extraction from 50 clinical specimens was performed using a commercial nucleic acid extraction kit (intron, Korea).

The extracted samples were examined at 260 nm and their purity was evaluated on A260-A280 basis. In this study, specific primers (HPV16 and HPV18) were used to identify HPV. The band size of the products on gel electrophoresis is mentioned in [Table 1](#).

Performing PCR to Identify HPV16 and HPV18 Genes

To perform multiplex polymerase chain reaction (PCR) in the final volume of 25 µl of master mix, 3 µl

of DNA with a 50 µl concentration, 10 µl of primer, and sterile distilled water were used according to the following protocol:

Primary denaturation at 95°C for 15 minutes followed by 40 seconds of secondary denaturation, annealing (primer bonding) at 58°C for 1 minute, initial expansion at 72°C for 2 minutes, and final expansion at 72°C for 7 minutes. Multiplex PCR products were electrophoresed on 1% agarose gel in 90 TBE XT buffer (Cinagen, Iran) for 90 minutes and analyzed using Gel documentation (9, 10).

Table 1. Primers used to propagate HPV 16, 18 (FR: forward primer, RP: reverse primer) 5

Primers	Primer sequence (5'-)	Target location	Product size
HPV-16	FP- 5'-TCA AAA GCC ACT GTG TCC TGA-3'	421-440	119 bp
	RP- 5'-CGT GTT CTT GAT GAT CTG CAA-3'	521-540	
HPV-18	FP- 5'-CCG AGC ACG ACA GGA ACG ACT-3'	533-553	172 bp
	RP- 5'-TCG TTT TCT TCC TCT GAG TCG CTT-3'	682-705	

FR: forward primer, RP: reverse primer

Results

This study aimed to determine the presence of HPVs in uterine cancer and the relationship between cancer morbidity and the virus in proportion to the virus type. Accordingly, samples were examined by molecular methods. [Table 2](#) shows the characteristics of the collected samples, along with specific PCR results of the two common types. Out of 50 patients with cervical cancer, 100% were HPV-positive. The

distribution of HPV types were as follows: 7 patients (14%) had HPV18, 19 (38%) had HPV16, 7 (14%) had HPV16 and HPV18, and 17 (34%) did not have any of the two types. In other words, the prevalence of HPV infection in this population equaled 66% ([Figure 1](#)). The positive control group had both types 16 and 18 while the negative control group had none.



Figure 1. Resulting band samples from PCR run on the agarose gel belonging to the HPV positive, negative and control samples.

+: Positive control / -: Negative control / 1: Does not have HPV16 or HPV18. / 2: Only HPV16 / 6: Only HPV18. / 5: Affected by both HPV16, 18.

Table 2. The prevalence of two common types of HPV among cervical cancer samples.

Total	HPV types						Pathological findings
	negative	positive	-16/18	+16/18	18	16	cervical cancer
50	17	33	17	7	7	19	Number of patients

Discussion

It has been proven that continuous HPV infection can lead to cervical cancer and intra epithelial neoplasia (11). Each year, more than 500,000 women are diagnosed with cervical cancer, mostly in developing countries (12,13). HPV has been found to be a necessary but not sufficient cause of cervical cancer (14). Serological methods and viral culture are not completely sensitive and reliable in terms of detecting HPV. Thus, application of precise and fast tests is gaining more popularity (15, 16).

In the study, the distribution of HPV types 16 and 18 was as follows: 7 patients (14%) had HPV18, 19 (38%) had HPV16, 7 (14%) had HPV16 and HPV18, and 17 patients (34%) did not have any of the two types.

More than 99% of uterine cancers are caused by HPV. European studies have mainly focused on HPV16 and HPV 18 variants (17).

In 2006, Meshkat *et al.* conducted a study on detecting provincial papillomavirus types 16 and 18 by restriction fragment length polymorphism (RFLP) PCR in cervical cancer samples isolated with paraffin. More than 200 types of HPV are known based on genomic DNA sequence, of which 85 types are well known and 120 types have new genotypes (18).

In 2016, Stamenkovic *et al.* examined the distribution of HPV in cervical cancer tissue. HPV genotype was identified in 19 to 22 positive sample. There were six high-risk genotypes, including 16, 18, 33, 45, 58, and 53 of which HPV16 was the most common (19).

In a study by Kan *et al.*, 50 samples of endocervical cancer were examined by PCR. HPV16, HPV18, HPV33, and HPV18 genomes were observed in two uterine cancer samples in Australian women. Out of 50 samples, 24 (48%) were HPV-positive (20).

In a study by Mahmoodi *et al.*, Which examined the prevalence of papilloma virus in uterine cancer, the results showed that 60 samples (43.3%) out of 72 samples were infected with HPV. The highest prevalence rate was observed in the age range of 48-63 years. HPV16 was the most prevalent papilloma virus (100%) (21).

In a study by Cocuzza *et al.*, which examined the prevalence of papillomavirus in uterine cancer samples, the results showed that 41 samples (34.2%) out of 120 plasmid samples were positive for papillomavirus DNA, and the prevalence of HPV45, HPV51, and HPV16 equaled 46.3%, 29.6%, and 18.5%, respectively (22).

Szostek *et al.* examined the physical state of HPV16 in cervical intraepithelial lesions and concluded that several factors contribute to the persistence and progression of HPV in cervical cancer, including suppression of the immune system due to infection with HPV or other microorganisms mentioned early onset of sexual life, long-term hormonal methods of contraception, smoking, and other sexually transmitted infections (23). In cervical cancer, due to the availability of relevant tissue, prevention, early diagnosis and timely treatment are effective in reducing mortality (24). In recent years, global scientific centers have increasingly focused on developing HPV vaccines (25). Due to the relatively high prevalence of the virus in cancer tissues, antiviral vaccines may be used in high-risk population in the future. Interferon drugs are also used to treat lesions associated with HPV infection (25).

Using this treatment is necessary to detect the presence of infection in the tissues. Therefore, it seems necessary to perform molecular tests along with other studies in precancerous and cervical cancer lesions and even suspicious cervicovaginal smears.

Conducting a cohort study in a larger population and in different provinces of Iran is suggested in order to accurately evaluate factors associated with acute leukemia, cancer risk factors and prevention.

Conclusion

Our study suggests that HPV16 was the most prevalent type (38%) in HPV-positive women. The study can also be beneficial for disease management and high sensitivity diagnosis of cervical intraepithelial neoplasia. Further studies should be conducted to check the HPV genotyping in other areas in Iran or other parts of the world.

Acknowledgments

This study was supported by by kerman university.

Conflict of Interest

The author declares that he has no conflicts of interest related to the subject matter or materials discussed in this article

References

- Muñoz N, Bosch FX, De Sanjosé S, Herrero R, Castellsagué X, Shah KV, et al. Epidemiologic classification of human papillomavirus types associated with cervical cancer. *N Engl J Med* 2003; 348(6): 518-27. [DOI:10.1056/NEJMoa021641] [PMID]
- Zinatizadeh MR, Masoumalinejad Z, Parnak F. Prevalence of Mycoplasma hyorhinis contamination in tissues samples from cancer patients: A Brief Report. *Mod Med Lab J* 2017; 1(3): 91-5 [DOI:10.30699/mmlj17.1.3.91]
- Izadimood N, Azadmanesh K, RahnamayeFarzami M. Development of In-House Multiplex Real Time PCR for Human Papillomavirus Genotyping in Iranian Women with Cervical Cancer and Cervical Intraepithelial Neoplasia. *Asian Pac J Cancer Prev* 2014;15(15):6257-6261. [DOI:10.7314/APJCP.2014.15.15.6257] [PMID]
- Sohrabi A, Mirab-Samiee S, Rahnamaye-Farzami M, Rafizadeh M, Akhavan S, Hashemi-Bahremani M, et al. C13orf18 And C1orf166 (MULAN) DNA Genes Methylation Are Not Associated with Cervical Cancer and Precancerous Lesions of Human Papillomavirus Genotypes In Iranian Women. *Asian Pac J Cancer Prev* 2014;15(16):6745-6748. [DOI:10.7314/APJCP.2014.15.16.6745] [PMID]
- Keshe MM, Kaffashi A, Bagheri Gh, Shahkarami MK, Mohammadi M, Nadji SA. [Identification of Human Papillomavirus Type 16 among Thinprep Samples from 11 Provinces of Iran. *IJOGI* 2013; 16(72): 22-2
- Shayanfar N, Hosseini N, Panahi M, Azadmanesh K, Mohammadpour M, Kadivar M, et al. Detection of Mucosal Type Human Papillomavirus in Cutaneous Squamous Cell Carcinoma in Iran. *Pathol Res Pract* 2013; 209: 90-94 [DOI:10.1016/j.prp.2012.10.010] [PMID]
- de Villiers EM. Cross-Roads in The Classification of Papillomaviruses. *Virology* 2013;445: 2-10. [DOI:10.1016/j.virol.2013.04.023] [PMID]
- Miller DL, Puricelli MD, Stack MS. Virology and Molecular Pathogenesis of HPV (Human Papillomavirus)-Associated Oropharyngeal Squamous Cell Carcinoma. *Biochem J* 2012; 443(2): 339-53. [DOI:10.1042/BJ20112017] [PMID] [PMCID]
- Peedicayil A. Human papillomavirus genotypes associated with cervical neoplasia in India. *Int J Gynecol Cancer*, 2006. 16(4). [DOI:10.1111/j.1525-1438.2006.00631.x] [PMID]
- Rajaram S. High-risk human papillomavirus, tumor suppressor protein p53 and mitomycin-C in invasive squamous cell carcinoma cervix. *Indian J Cancer*, 2006. 43(4): p. 156. [DOI:10.4103/0019-509X.29420] [PMID]
- Dorostkar R. Molecular Recognition of Human Papilloma Virus (HPV) Using Proprietary PCR Method Based on L1 Gene and the Evaluation of its Frequency in Tissue Samples from Patients with Cervical Cancer. *Arak Uni Med Sci J*, 2015. 18(3): 28-36.
- Poljak M. Human papillomavirus prevalence and type-distribution, cervical cancer screening practices and current status of vaccination implementation in Central and Eastern Europe. *Vaccine*, 2013. 31: H59-H70. [DOI:10.1016/j.vaccine.2013.03.029] [PMID]
- Hassani M, Salehian P, Pourazar S. Human Papiloma Virus detection in various cervical lesions by molecular methods. *Sarem J Reprod Med*, 2017. 1(3):113-116. [DOI:10.29252/sjrm.1.3.113]
- You K. High-risk human papillomavirus DNA testing and high-grade cervical intraepithelial lesions. *Aust N Z J Obstet Gynaecol*, 2007. 47(2):141-144. [DOI:10.1111/j.1479-828X.2007.00701.x] [PMID]
- Loya A. Human Papillomavirus genotype distribution in invasive cervical cancer in Pakistan. *Cancers*, 2016. 8(8):72. [DOI:10.3390/cancers8080072] [PMID] [PMCID]
- Ouédraogo C. Distribution of high-risk human papillomavirus genotypes in precancerous cervical lesions in Ouagadougou, Burkina Faso. *Open J Obstet Gynecol*, 2016. 6(4): 196-204. [DOI:10.4236/ojog.2016.64025]
- Schiller JT, Hidesheim A. Developing HPV virus-like particle vaccines to prevent cervical cancer: a progress report. *J Clin Virol*. 2000;19(1-2):67-74. [DOI:10.1016/S1386-6532(00)00091-3]
- Meshkat Z, Soleimanjahi H, Mahmoudi M, Mirshahabi H, Hassan ZM, Ghaffari SR, et al. Determination of human papillomavirus type 16 genotype and construction of cloning vector pTZ57R encoding HPV16 E7 gene. *Saudi Med J*. 2007;28(10):1511-5.
- Stamenkovic M, Knezevic A, Knezevic I, Kuzmanovic I, Karalic D, Milenkovic S, et al. Highrisk human papilloma virus genotypes in cervical carcinoma of Serbian women: Distribution and association with pathohistological findings. *Biologicals*. 2016;44(5):412-6. [DOI:10.1016/j.biologicals.2016.05.001] [PMID]
- Kan CY, Iacopetta BJ, Lawson JS, Whitaker NJ. Identification of human papillomavirus DNA gene sequences in human breast cancer. *Br J Cancer*. 2005;93(8):946-8. [DOI:10.1038/sj.bjc.6602778] [PMID] [PMCID]
- Mahmoodi P, Motamedi H, Reza Seyfi Abad Shapouri M, Bahrami Shehni M, and Kargar M. Molecular Detection and Typing of Human Papillomaviruses in Paraffin-Embedded Cervical Cancer and Pre-Cancer Tissue Specimens. *Iran J Cancer Prev*. 2016; 9(1): e3752. [DOI:10.17795/ijcp-3752] [PMID] [PMCID]
- Cocuzza CE, Martinelli M, Sina F, Piana A, Sotgiu G, Dell Anna T, Musumeci R. Human papillomavirus DNA detection in plasma and cervical samples of women with a recent history of low grade or precancerous cervical dysplasia. *PLoS One*. 2017; 12(11): e0188592. [DOI:10.1371/journal.pone.0188592] [PMID] [PMCID]
- Szostek S, Zawilinska B, Kopec J, KoszVnenchak M. Herpesviruses as possible cofactors in HPV-16-related oncogenesis. *Acta Biochim Pol* 2009; 56(2): 337. [DOI:10.18388/abp.2009_2466] [PMID]

24. McNair R, Power J, Carr S. Comparing knowledge and perceived risk related to the human papilloma virus among Australian women of diverse sexual orientations. *Aust N Z J Public Health*. 2009;33(1):87-93. [[DOI:10.1111/j.1753-6405.2009.00345.x](https://doi.org/10.1111/j.1753-6405.2009.00345.x)] [[PMID](#)]
25. Keyhani E, Kohannia N, Izadimood N, Keykhaee M, Najmabadi H. The prevalence of human papilloma virus (HPV) in malignant cervical lesion, using multiplex PCR. *Tehran Univ Med J TUMS Publ*. 2006;64(3):95

How to Cite This Article:

Shahi Z, Edalatmanesh M A, kheirkhah B. Molecular Detection of Human Papilloma Virus (Type 16,18) Using PCR and Its Frequency in Patients with Cervical Cancer in Iranian Women. *J Obstet Gynecol Cancer Res*. 2020; 5 (3) :110-114

Download citation:

[BibTeX](#) | [RIS](#) | [EndNote](#) | [Medlars](#) | [ProCite](#) | [Reference Manager](#) | [RefWorks](#)